Functional Ecology 2002
16, 870–873

Author Index

Addo-Bediako, A., Chown, S. L. & Gaston, K. J. Metabolic cold adaptation in insects: a large-scale perspective, 332
Aerts, P. *see* Berwaerts, K.
Ågren G. I. *see* Franklin O.
Aguilera, A. *see* Rey, P. J.
Alcántara, J. M. *see* Rey, P. J.
Alexander, I. J. *see* Genney, D. R.
Alfaro, M. E. Forward attack modes of aquatic feeding garter snakes, 204
Allison V. J. Nutrients, arbuscular mycorrhizas and competition interact to influence seed production and germination success in *Achillea millefolium*, 742
Allison, V. J. & Goldberg, D. E. Species-level versus community-level patterns of mycorrhizal dependence on phosphorus: an example of Simpson's paradox, 346
Almeida-Cortez, J. S. & Shipley, W. No significant relationship exists between seedling relative growth rate under nutrient limitation and potential tissue toxicity, 122
Alpert, P., Holzapfel, C. & Benson, J. M. Hormonal modification of resource sharing in the clonal plant *Fragaria chiloensis*, 191
Álvarez, D. & Nicieza, A. G. Effects of temperature and food quality on anuran larval growth and metamorphosis, 640
Anthony, P. A., Holtum J. A. M. & Jackes B. R. Shade acclimation of rainforest leaves to colonization by lichens, 808

Baeza, J. A. & Fernández, M. Active brood care in *Cancer setosus* (Crustacea: Decapoda): the relationship between female behaviour, embryo oxygen consumption and the cost of brooding, 241
Barclay, R. M. R. *see* Chruszcz, B. J.
Barker, G. M. *see* Wardle, D. A.
Barker, R. J. *see* Hall, A. J.
Barnes D. K. A. & De Grave S. Temporospatial constraints in resources available to and used by hermit crabs: tests of models, 714
Baur, B. *see* Locher, R.
Bellamy P. E. *see* Hinsley, S. A.
Bellwood, D. R., Wainwright, P. C., Fulton, C. J. & Hoey, A. Assembly rules and functional groups at global biogeographical scales, 557
Benson, J. M. *see* Alpert, P.
Berendse, F. *see* van der Krift, T. A. J.
Berwaerts, K., Van Dyck, H. & Aerts, P. Does flight morphology relate to flight performance? An experimental test with the butterfly *Pararge aegeria*, 484
Bevan, R. M., Butler, P. J., Woakes, A. J. & Boyd, I. L. The energetics of Gentoo Penguins, *Pygoscelis papua*, during the breeding season, 175
Bin Hashim, R. *see* Heil, M.
Blount, J. D., Surai, P. F., Houston, D. C. & Møller, A. P. Patterns of yolk enrichment with dietary carotenoids in gulls: the roles of pigment acquisition and utilization, 445
Boddy, L. *see* Wells, J. M.
Bonner, K. I. *see* Wardle, D. A.
Borghetti, M. *see* Magnani, F.
Boyd, I. L. *see* Bevan, R. M.
Boyd, I. L. *see* Green, J. A.
Bragg, J. G. & Westoby, M. Leaf size and foraging for light in a sclerophyll woodland, 633
Brooks, J. R. *see* Buchmann, N.
Bryant D. M. *see* Engstrand S. M.
Buchmann, N., Brooks, J. R. & Ehleringer, J. R. Predicting daytime carbon isotope ratios of atmospheric CO_2 within forest canopies, 49
Buck, T. L. *see* Zimmer, M.
Burslem, D. F. R. P. *see* Daws, M. I.
Bustnes, J. O., Folstad, I., Erikstad, K. E., Fjeld, M., Miland, Ø. O. & Skaare, J. U. Blood concentration of organochlorine pollutants and wing feather asymmetry in Glaucous Gulls, 617
Butler, P. J. *see* Bevan, R. M.
Butler, P. J. *see* Green, J. A.

Callaway R. M. *see* Zabinski, C. A.
Carefoot, T. H. *see* Zimmer, M.
Carey, J. R., Liedo, P., Harshman, L., Liu, X., Müller, H.-G., Partridge, L. & Wang, J.-L. Food pulses increase longevity and induce cyclical egg production in Mediterranean fruit flies, 313
Casas J. *see* Giron, D.
Chapple D. G. & Swain R. Effect of caudal autotomy on locomotor performance in a viviparous skink, *Niveoscincus metallicus*, 817
Chown, S. L. *see* Addo-Bediako, A.
Chrictian, R. *see* Turrnbull, M. H.
Chruszcz, B. J. & Barclay, R. M. R. Thermoregulatory ecology of a solitary bat, *Myotis evotis*, roosting in rock crevices, 18
Coners, H. & Leuschner, C. *In situ* water absorption by tree fine roots measured in real time using miniature sap-flow gauges, 696
Coulson, S. J., Hodkinson, I. D., Webb, N. R. & Harrison, J. A. Survival of terrestrial soil-dwelling arthropods on and in seawater: implications for trans-oceanic dispersal, 353
Cowling, R. M. *see* Lamont, B. B.
Crabtree, L. M. *see* Daws, M. I.
Craine, J. M., Tilman, D., Wedin, D., Reich, P., Tjoelker, M. & Knops, J. Functional traits, productivity and effects on nitrogen cycling of 33 grassland species, 563
Cresswell, W. *see* Reid, J. M.
Cutts, C. J., Metcalfe, N. B. & Taylor, A. C. Juvenile Atlantic Salmon (*Salmo salar*) with relatively high standard metabolic rates have small metabolic scopes, 74

Dalling, J. W. *see* Daws, M. I.
Darrouzet E. *see* Giron, D.
Daws, M. I., Burslem, D. F. R. P., Crabtree, L. M., Kirkman, P., Mullins, C. E. & Dalling, J. W. Differences in seed germination responses may promote coexistence of four sympatric *Piper* species, 258
De Grave S. *see* Barnes D. K. A.
Delucia, E. H. *see* Maherali, H.
Digneffe, G. *see* Pakeman, R. J.
Dohm, M. R. Repeatability estimates do not always set an upper limit to heritability, 273
Doncaster, C. P. *see* Rondinini, C.
Donovan, L. A. *see* Espeleta, J. F.
Dore, S. *see* Reichstein, M.
Dormann, C. F. & Skarpe, C. Flowering, growth and defence in the two sexes: consequences of herbivore exclusion for *Salix polaris*, 649
Dormann, C. F. & Woodin, S. J. Climate change in the Arctic: using plant functional types in a meta-analysis of field experiments, 4
Douglas, A. E. *see* Johnson, S. N.
van Dyck, H. *see* Berwaerts, K.

Ehleringer, J. R. *see* Buchmann, N.
Eissenstat, D. M. *see* Yanai, R. D.
Elle, E. & Hare, J. D. Environmentally induced variation in floral traits affects the mating system in *Datura wrightii*, 79
Engstrand S. M. & Bryant D. M. A trade-off between clutch size and incubation efficiency in the Barn Swallow *Hirundo rustica*, 782

© 2002 British Ecological Society

Erikstad, K. E. *see* Bustnes, J. O.
Espeleta, J. F. & Donovan, L. A. Fine root demography and morphology in response to soil resource availability among xeric and mesic sandhill tree species, 113

Fernández, M. *see* Baeza, J. A.
Fernández, R. *see* Rey, P. J.
Fiala, B. *see* Heil, M.
Fjeld, M. *see* Bustnes, J. O.
Folstad, I. *see* Bustnes, J. O.
Franklin O. & Ågren G. I. Leaf senescence and resorption as mechanisms of maximizing photosynthetic production during canopy development at N limitation, 727
Fritz, H. *see* Guillemain, M.
Fulton, C. J. *see* Bellwood, D. R.

García, L. *see* Rey, P. J.
Garnier, E. *see* Lavorel, S.
Garrido, J. L. *see* Rey, P. J.
Gaston, K. J. *see* Addo-Bediako, A.
Gaveau D. L. A. *see* Hinsley, S. A.
Gawelczyk, A. T. *see* Kozłowski, J.
Genney, D. R., Alexander, I. J. & Hartley, S. E. Soil organic matter distribution and below-ground competition between *Calluna vulgaris* and *Nardus stricta*, 664
Giron, D., Rivero, A., Mandon, N., Darrouzet E. & Casas J. The physiology of host feeding in parasitic wasps: implications for survival, 750
Gleeson, T. T. *see* Hancock, T. V.
Goldberg, D. E. *see* Allison, V. J.
Goss-Custard, J. D. *see* Nagarajan, R.
Grace, J. *see* Magnani, F.
Green, J. A., Butler, P. J., Woakes, A. J. & Boyd, I. L. Energy requirements of female Macaroni Penguins breeding at South Georgia, 671
Groom, P. K. *see* Lamont, B. B.
Groothuis, T. G. & Schwabl, H. Determinants of within- and among-clutch variation in levels of maternal hormones in Black-Headed Gull eggs, 281
Grubb, P. J. *see* Metcalfe, D. J.
Guillemain, M., Martin, G. R. & Fritz, H. Feeding methods, visual fields and vigilance in dabbling ducks (Anatidae), 522

Hacke, U. G. *see* Sperry, J. S.
Hall, A. J., McConnell, B. J. & Barker, R. J. The effect of total immunoglobulin levels, mass and condition on the first-year survival of Grey Seal pups, 462
Hancock, T. V. & Gleeson, T. T. Metabolic recovery in the Desert Iguana (*Dipsosaurus dorsalis*) following activities of varied intensity and duration, 40
Hanley, M. E. & Lamont, B. B. Relationships between physical and chemical attributes of congeneric seedlings: how important is seedling defence? 216
Hare, J. D. *see* Elle, E.
Harrison, J. A. *see* Coulson, S. J.
Harshman, L. *see* Carey, J. R.
Hartley, F. G. *see* Iason, G. R.
Hartley, S. E. *see* Genney, D. R.
Hartley, S. E. *see* Johnson, S. N.
Heil, M., Hilpert, A., Fiala, B., Bin Hashim, R., Strohm, E., Zotz, G. & Linsenmair, K. E. Nutrient allocation of *Macaranga triloba* ant plants to growth, photosynthesis and indirect defence, 475
Herrel A. *see* Verwaijen, D.
Hikosaka, K. *see* Yasumura, Y.
Hill, R. A. *see* Hinsley, S. A.
Hilpert, A. *see* Heil, M.
Hinsley, S. A., Hill, R. A. Gaveau D. L. A. & Bellamy P. E. Quantifying woodland structure and habitat quality for birds using airborne laser scanning, 851
Hirose T. *see* Yasumura, Y.
Hodkinson, I. D. *see* Coulson, S. J.
Hoey, A. *see* Bellwood, D. R.
Högberg, P. *see* Jumpponen, A.
Högstedt, G. *see* Paulsen, T. R.
Holt, S. *see* Reid, J. M.
Holtum J. A. M. *see* Anthony, P. A.
Holzapfel, C. *see* Alpert, P.
Houston, D. C. *see* Blount, J. D.
Howell, C. J., Kelly, D. & Turnbull, M. H. Moa ghosts exorcised? New Zealand's divaricate shrubs avoid photoinhibition, 232
Howell, C. T. *see* Turnbull M. H.
Hulme, P. E. *see* Willis, S. G.
Huss-Danell, K. *see* Jumpponen, A.
Huuskonen, H. *see* Lahti, K.

Iason, G. R., Manso, T., Sim, D. A. & Hartley, F. G. The functional response does not predict the local distribution of European Rabbits (*Oryctolagus cuniculus*) on grass swards: experimental evidence, 394
Isebrands J. G. *see* McDonald, E. P.

Jackes B. R. *see* Anthony, P. A.
Jackson, S. & Wilson, R. P. The potential costs of flipperbands to penguins, 141
Johnson, S. N., Mayhew, P. J., Douglas, A. E. & Hartley, S. E. Insects as leaf engineers: can leaf-miners alter leaf structure for birch aphids? 575
Jumpponen, A., Högberg, P., Huss-Danell, K. & Mulder, C. P. H. Interspecific and spatial differences in nitrogen uptake in monocultures and two-species mixtures in north European grasslands, 454

Kelly, D. *see* Howell, C. J.
Kelly, D. *see* Turnbull, M. H.
King R. B. Predicted and observed maximum prey size – snake size allometry, 766
Kirkman, P. *see* Daws, M. I.
Kitajima, K. Do shade-tolerant tropical tree seedlings depend longer on seed reserves? Functional growth analysis of three Bignoniaceae species, 433
Knops, J. *see* Craine, J. M.
Kozłowski, J. Theoretical and empirical status of Brown, Marquet and Taper's model of species–size distribution, 540
Kozłowski, J. & Gawelczyk, A. T. Why are species' body size distributions usually skewed to the right? 419
van der Krift, T. A. J. & Berendse, F. Root life spans of four grass species from habitats differing in nutrient availability, 198
Kruger, E. L. *see* McDonald, E. P.

Ladley, J. J. *see* Norton, D. A.
Lahti, K., Huuskonen, H., Laurila, A. & Piironen, J. Metabolic rate and aggressiveness between Brown Trout populations, 167
Lamont, B. B., Groom, P. K. & Cowling, R. M. High leaf mass per area of related species assemblages may reflect low rainfall and carbon isotope discrimination rather than low phosphorus and nitrogen concentrations, 403
Lamont, B. B. *see* Hanley, M. E.
Laurila, A. *see* Lahti, K.
Lavorel, S. & Garnier, E. Predicting changes in community composition and ecosystem functioning from plant traits: revisiting the Holy Grail, 545
Lea, S. E. G. *see* Nagarajan, R.
Leuschner, C. *see* Coners, H.
Liedo, P. *see* Carey, J. R.
Liggins, G. A. *see* Lovvorn, J. R.
Linder H. P. *see* Verboom, G. A.
Linsenmair, K. E. *see* Heil, M.
Liu, X. *see* Carey, J. R.
Locher, R. & Baur, B. Nutritional stress changes sex-specific reproductive allocation in the simultaneously hermaphroditic land snail *Arianta arbustorum*, 623
Loeschcke, V. *see* Sørensen, J. G.

© 2002 British Ecological Society,

Loudon, C. & Zhang, J. Measuring drag without a force transducer: a terminal velocity assay, 268
Lovvorn, J. R. & Liggins, G. A. Interactions of body shape, body size and stroke-acceleration patterns in costs of underwater swimming by birds, 106
Lusk, C. H. Does photoinhibition avoidance explain divarication in the New Zealand flora? 858

Magnani, F., Grace, J. & Borghetti, M. Adjustment of tree structure in response to the environment under hydraulic constraints, 385
Maherali, H., Williams, B. L., Paige, K. N. & Delucia, E. H. Hydraulic differentiation of Ponderosa pine populations along a climate gradient is not associated with ecotypic divergence, 510
Malo, J. E. Modelling unimodal flowering phenology with exponential sine equations, 413
Mandon, N. *see* Giron, D.
Manso, T. *see* Iason, G. R.
Manzaneda A. J. *see* Rey, P. J.
Martin, G. R. *see* Guillemain, M.
Matsui K. *see* Yasumura, Y.
Mayhew, P. J. *see* Johnson, S. N.
McConnell, B. J. *see* Hall, A. J.
McDonald, E. P., Kruger, E. L., Riemenschneider D. E. & Isebrands J. G. Competitive status influences tree-growth responses to elevated CO_2 and O_3 in aggrading aspen stands, 792
McLain, D. K. *see* Pratt, A. E.
Mellanby, R. J. *see* Reid, J. M.
Metcalfe, D. J., Grubb, P. J. & Metcalfe, S. S. Soil dilution as a surrogate for root competition: effects on growth of seedlings of Australian tropical rainforest trees, 223
Metcalfe, N. B. *see* Cutts, C. J.
Metcalfe, S. S. *see* Metcalfe, D. J.
Meziane, D. *see* Shipley, B.
Miland, Ø. O. *see* Bustnes, J. O.
Møller, A. P. *see see* Blount, J. D.
Moya-Laraño J. Senescence and food limitation in a slowly ageing spider, 734
Mulder, C. P. H. *see* Jumpponen, A.
Müller, H.-G. *see* Carey, J. R.
Mullins, C. E. *see* Daws, M. I.

Nagarajan, R., Lea, S. E. G. & Goss-Custard, J. D. Mussel valve discrimination and strategies used in valve discrimination by the Oystercatcher, *Haematopus ostralegus*, 339
Nakamura, T., Uemura, S. & Yabe, K. Variation in nitrogen-use traits within and between five *Carex* species growing in the lowland mires of northern Japan, 67
Nentwig W. *see* Wullschleger B.
Nicieza, A. G. *see* Álvarez, D.
Niklaus, P. A. *see* Volk, M.
Norton, D. A., Ladley, J. J. & Sparrow, A. D. Host provenance effects on germination and establishment of two New Zealand mistletoes (Loranthaceae), 657

Ohashi, K. & Yahara, T. Visit larger displays but probe proportionally fewer flowers: counterintuitive behaviour of nectar-collecting bumble bees achieves an ideal free distribution, 492
Olofsson, C. *see* Olsson, M.
Olsson, M., Wapstra, E. & Olofsson, C. Offspring size–number strategies: experimental manipulation of offspring size in a viviparous lizard (*Lacerta vivipara*), 135
Ourcival, J.-M. *see* Reichstein, M.

Paige, K. N. *see* Maherali, H.
Pakeman, R. J., Digneffe, G. & Small, J. L. Ecological correlates of endozoochory by herbivores, 296
Pannell, J. R. What is functional androdioecy? 862
Partridge, L. *see* Carey, J. R.
Passy, S. I. Environmental randomness underlies morphological complexity of colonial diatoms, 690
Paulsen, T. R. & Högstedt, G. Passage through bird guts increases germination rate and seedling growth in *Sorbus aucuparia*, 608
Pennings, S. C. *see* Zimmer, M.
Piironen, J. *see* Lahti, K.
Plaut, I. Does pregnancy affect swimming performance of female Mosquitofish, *Gambusia affinis*? 290
Pratt, A. E. & McLain, D. K. Antisymmetry in male fiddler crabs and the decision to feed or breed, 89

Quinn L. *see* Zabinski, C. A.

Rambal, S. *see* Reichstein, M.
Ramírez, J. M. *see* Rey, P. J.
Reich, P. *see* Craine, J. M.
Reichstein, M., Tenhunen, J. D., Roupsard, O., Ourcival, J.-M., Rambal, S., Dore, S. & Valentini, R. Ecosystem respiration in two Mediterranean evergreen Holm Oak forests: drought effects and decomposition dynamics, 27
Reid, J. M., Cresswell, W., Holt, S., Mellanby, R. J., Whitfield, D. P. & Ruxton, G. D. Nest scrape design and clutch heat loss in Pectoral Sandpipers (*Calidris melanotos*), 305
Rey, P. J., Garrido, J. L., Alcántara, J. M., Ramírez, J. M., Aguilera, A., García, L., Manzaneda A. J. & Fernández R. Spatial variation in ant and rodent post-dispersal predation of vertebrate-dispersed seeds, 773
Riemenschneider D. E. *see* McDonald, E. P.
Rivero, A. *see* Giron, D.
Robinson, D. *see* Gaston, K.
Rödl, T. & Ward, D. Host recognition in a desert mistletoe: early stages of development are influenced by substrate and host origin, 128
Rondinini, C. & Doncaster, C. P. Roads as barriers to movement for hedgehogs, 504
Roupsard, O. *see* Reichstein, M.
Ruiz, N., Ward, D. & Saltz, D. Calcium oxalate crystals in leaves of *Pancratium sickenbergeri*: constitutive or induced defence? 99
Rukke, N. A. Effects of low calcium concentrations on two common freshwater crustaceans, *Gammarus lacustris* and *Astacus astacus*, 357
Ruxton, G. D. *see* Reid, J. M.

Saltz, D. *see* Ruiz, N.
Schwabl, H. *see* Groothuis, T. G.
Shipley, B. Trade-offs between net assimilation rate and specific leaf area in determining relative growth rate: relationship with daily irradiance, 682
Shipley, B. & Meziane, D. The balanced-growth hypothesis and the allometry of leaf and root biomass allocation, 326
Shipley, W. *see* Almeida-Cortez, J. S.
Sim, D. A. *see* Iason, G. R.
Skaare, J. U. *see* Bustnes, J. O.
Skarpe, C. *see* Dormann, C. F.
Small, J. L. *see* Pakeman, R. J.
Sørensen, J. G. & Loeschcke, V. Decreased heat-shock resistance and down-regulation of Hsp70 expression with increasing age in adult *Drosophila melanogaster*, 379
Sparrow, A. D. *see* Norton, D. A.
Sperry, J. S. & Hacke, U. G. Desert shrub water relations with respect to soil characteristics and plant functional type, 367
Stelzer C. P. Phenotypic plasticity of body size at different temperatures in a planktonic rotifer: mechanisms and adaptive significance, 835
Stock W. D. *see* Verboom, G. A.
Strohm, E. *see* Heil, M.
Surai, P. F. *see* Blount, J. D.
Swain R. *see* Chapple D. G.

© 2002 British Ecological Society,

Takakura, K. The specialist seed predator *Bruchidius dorsalis* (Coleoptera: Bruchidae) plays a crucial role in the seed germination of its host plant, *Gleditsia japonica* (Leguminosae), 252
Taylor, A. C. *see* Cutts, C. J.
Tenhunen, J. D. *see* Reichstein, M.
Tilman, D. *see* Craine, J. M.
Tjoelker, M. *see* Craine, J. M.
Turnbull, M. H. *see* Howell, C. J.
Turnbull, M. H., Howell, C. J., Christian, R. & Kelly, D. Photoinhibition, acclimation and New Zealand's divaricate plants: a reply to Lusk, 860

Uemura, S. *see* Nakamura, T.

Valentini, R. *see* Reichstein, M.
Van Damme R. *see* Verwaijen, D.
Verboom, G. A., Stock W. D. & Linder H. P. Determinants of postfire flowering in the geophytic grass *Ehrharta capensis*, 705
Verwaijen, D., Van Damme R. & Herrel A. Relationships between head size, bite force, prey handling efficiency and diet in two sympatric lacertid lizards, 842
Volk, M. & Niklaus, P. A. Respiratory carbon loss of calcareous grasslands in winter shows no effects of 4 years' CO_2 enrichment, 162

Wainwright, P. C. *see* Bellwood, D. R.
Wang, J.-L. *see* Carey, J. R.
Wapstra, E. *see* Olsson, M.
Ward, D. *see* Rödl, T.
Ward, D. *see* Ruiz, N.
Wardle, D. A., Bonner, K. I. & Barker, G. M. Linkages between plant litter decomposition, litter quality, and vegetation responses to herbivores, 585
Webb, N. R. *see* Coulson, S. J.
Wedin, D. *see* Craine, J. M.
Wells, J. M. & Boddy, L. Interspecific carbon exchange and cost of interactions between basidiomycete mycelia in soil and wood, 153
Westoby, M. *see* Bragg, J. G.
Whitfield, D. P. *see* Reid, J. M.
Williams, B. L. *see* Maherali, H.
Willis, S. G. & Hulme, P. E. Does temperature limit the invasion of *Impatiens glandulifera* and *Heracleum mantegazzianum* in the UK? 530
Wilson, R. P. *see* Jackson, S.
Woakes, A. J. *see* Bevan, R. M.
Woakes, A. J. *see* Green, J. A.
Woodin, S. J. *see* Dormann, C. F.
Wullschleger B. & Nentwig W. Influence of venom availability on a spider's prey-choice behaviour, 802
Wunder, B. A. *see* Zynel, C. Y.

Yabe, K. *see* Nakamura, T.
Yahara, T. *see* Ohashi, K.
Yanai, R. D. & Eillenstat, D. M. Coping with herbivores and pathogens: a model of optimal root turnover, 865
Yasumura, Y., Hikosaka, K., Matsui K. & Hirose T. Leaf-level nitrogen-use efficiency of canopy and understorey species in a beech forest, 826

Zabinski, C. A., Quinn L. & Callaway R. M. Phosphorus uptake, not carbon transfer, explains arbuscular mycorrhizal enhancement of *Centaurea maculosa* in the presence of native grassland species, 758
Zhang, J. *see* Loudon, C.
Zimmer, M., Pennings, S. C., Buck, T. L. & Carefoot, T. H. Species-specific patterns of litter processing by terrestrial isopods (Isopoda: Oniscidea) in high intertidal salt marshes and coastal forests, 596
Zotz, G. *see* Heil, M.
Zynel, C. Y. & Wunder, B. A. Limits to food intake by the Prairie Vole: effects of time for digestion, 58

© 2002 British
Ecological Society,

Functional Ecology 2002
16, 874–877

Subject Index

absorptance spectra 808–816
Acacia raddiana 128–134
Achillea millefolium 454–461, 742–749, 758–765
Adrisa magna 252–257
age and size 640–648
ageing 379–384
aggression 73–78, 167–174
agonistic trout 167–174
Alepis flavida 657–663
algae 690–695
allocation 162–166, 326–331, 664–670
  rules 750–757
allometric engineering 135–140
allometry 326–331, 766–772
Amaranthus cruentus 727–733
Anas spp. 522–529
ant seed predation 773–781
ant–plant mutualism 475–483
antioxidants 445–453
antipredator 394–402
antisymmetry 89–98
anuran larvae 640–648
Apodemus speciosus 252–257
aphid 575–584
Aptenodytes patagonicus 141–148
arctic 4–17, 617–622, 649–656
Arianta arbustorum 623–632
Aristotelia fruticosa 232–240
Arrhenatherum elatius 198–203
arthropods 353–356
assembly rules 557–562
Astacus astacus 357–366
Asteraceae 122–127
asymmetry 339–345, 617–622
asynchrony 281–289
Atriplex spp. 367–378
attack strategies 204–215
avian gut treatment 608–616

balanced growth 326–331
Barents Sea 353–356
basidiomycete mycelia 153–161
beech 696–703, 826–834
beetle 252–257
behaviour 273–280, 492–503, 504–509
  optimal construction 305–312
Bergmann's rule 835–841
Betula pendula 575–584
Bignoniaceae 433–444
biodiversity 557–562, 596–607
biogeochemical cycles 545–556
biomass 4–17, 326–331, 742–749
biophysics 268–272
birds 106–112, 305–312, 608–616, 851–857
  ducks 522–529
  gulls 281–289, 445–453, 617–622
  oystercatcher 339–345
  penguins 141–148, 175–190, 671–681
  swallow 782–791
bite force 842–850
Blatta orientalis 802–807
Bombus diversus 492–503
breeding season 175–190, 671–681
brood care 175–190, 241–251, 671–681
brood reduction 281–289
brooding female 241–251
Bruchidius dorsalis 252–257

calcareous grassland 162–166
calcification 357–366
calibration 696–703
Calidris melanotos 305–13
Calluna vulgaris 664–70
Camisia anomia 353–356
Cancer setosus 241–251
canopy 49–57, 232–240, 585–595, 727–733, 851–857
carbon 27–39, 153–161, 162–166, 232–240
  transfer 191–197, 758–765
carbon dioxide 162–166, 758–765, 792–801
  exchange 808–816
carbon isotope discrimination 49–57, 403–412
Carex lyngbyei 67–72
carotenoid metabolism 445–453
Cassiope tetragona 4–17
Centaurea maculosa 758–765
Challichlamys latifolia 433–444
cheliped 89–98
Chiroptera 18–26
chlorophyll 232–240, 808–816
Chrysothamnus spp. 367–378
Cirsium purpuratum 492–503
Cistus spp. 413–418
climate 4–17, 385–393, 510–521
  change 530–539, 792–801
clonal plants 191–197, 792–801
Coleoptera 252–257
Collembola 353–356
colonization 353–356, 530–539, 808–816
community composition 346–352, 545–556
competition 623–632, 664–670, 714–726, 792–801
Coprosma propinqua 232–240
coral reef 557–62
Corokia cotoneaster 232–40
costs 289–295, 305–312, 782–791, 817–825
  swimming 106–112, 141–148, 175–190
crabs 241–251, 714–720
critical swimming speed 290–295
Crustacea 89–98, 241–251, 353–356, 596–607, 714–720
  freshwater 357–366
Cupiennius salei 802–807

δ^{13} 49–57
dabbling ducks 522–529
Datura wrightii 79–88
decay 153–161, 585–595, 596–607
decomposition 585–595, 596–607
  dynamics model 27–39
defence 99–105, 232–240, 475–483
  chemical 122–127, 216–222, 574–584, 649–656
defoliation 705–713
desert 40–48, 367–378, 510–521
  Negev 99–105, 128–135
diatoms 690–695
digestion 58–66, 596–607
Dipsosaurus dorsalis 40–48
Diploglottis bracteata 223–231
Discoglossus galganoi 640–648
discriminant function analysis 296–304
dispersal 296–304, 353–356, 773–781
disturbance 545–556
dominance 73–78
Drosophila melanogaster 379–384
drought 27–39

ecological barrier 504–509
ecological immunology 462–474
ecosystem functioning 545–556
ecosystem respiration 27–39
ecotypes 510–521
eddy covariance 27–39

© 2002 British Ecological Society

effect traits 545–556
eggs 135–140, 281–289, 313–325, 835–841
  incubation costs 305–312
Ehrharta capensis 705–713
Elaeocarpus grandis 223–231
embryo mass 241–251
endozoochory 296–304
energetics 40–48, 73–78
  penguins 141–148, 175–190, 671–681
  swimming costs 106–112, 141–148
energy 58–66, 433–444
environmental effects 273–280
environmental gradients 346–352
Erinaceus europaeus 504–509
Eriocrania spp. 575–584
establishment 657–663
Euceraphis betulae 575–584
Eudyptes spp. 141–148, 671–681
Eupelmus vuilletti 750–757
Euphausia superba 671–681
evolutionary strategy 690–695
extinction 419–432

Fagus spp. 696–703, 826–834
feeding bouts 58–66
Festuca spp. 394–402, 454–461, 758–765
fire 545–556, 705–713
fish 73–78, 167–174, 290–295, 557–562
fitness 742–749
flight 484–491
flightlessness 332–338
flipper-bands 141–148
floral display 79–88, 492–503, 705–713
flower revisiting 492–503
flowering phenology model 413–418
fluctuating asymmetry 89–98, 617–622
fluid drag model 268–272
foliage 49–57
Folsomia quadrioculata 353–356
food restriction 313–325, 734–741
foot propulsion 106–112
foraging 522–529, 633–639, 750–757
  behaviour 339–345, 394–402, 492–503
  costs 141–148
  penguin 175–190, 671–681
forest 27–39, 585–595, 596–607, 792–801
  beech 696–703, 826–34
  canopy 49–57, 585–595, 727–733, 826–834
  tropics 223–231, 433–444, 808–816
    niche species 258–267
fractal dimension 690–695
Fragaria chiloensis 191–197
frogs 640–648
frugivory 608–616
fruit flies 313–325
functional classifications 563–574
functional groups 557–562
functional morphology 204–215
functional response 394–402
fynbos 705–713

Gaillardia aristata 758–765
Gambusia affinis 290–295
Gammarus lacustris 357–366
gape in snakes 204–215, 766–772
Gazella dorcas 99–105
genetic variation 510–521
genotype–environment interaction 273–280
germination 252–257, 258–267, 657–663
  rate 608–616
Gleditsia japonica 252–257
global change 4–17, 792–801
Glycine max 727–733
gradients 346–352
grass sward 394–402
grasses 4–17, 198–203, 454–461, 705–713
grassland 162–166, 563–574, 758–765
Grayia spinosa 367–378
growth 357–366, 640–648, 792–801
  analysis 433–444
  differentiation balance 475–483
gut survival 296–304, 608–618

habitat 67–72, 394–402, 851–857
Haematopus ostralegus 339–345
Hakea spp. 216–222, 403–412
Halichoerus grypus 462–474
Halimium sp. 413–418
hammering shells 339–345
hard seed germination 252–257
hatching asynchrony 281–289
heart rate 175–190, 671–681
heat resistance 379–384
heat-balance approach 696–703
heathland 664–670
Heracleum mantegazzianum 530–539
herbaceous allocation 326–331
herbivores 296–304, 585–595, 649–656
herbivory 323–40, 394–402, 649–56
  defence 99–105, 216–222
  perennial herb 79–88
heritability 273–280
herkogamy 79–88
hermaphrodite 623–632
Hirundo rustica 782–791
holdfast formation 128–134
Holm Oak 27–39
homeostasis 385–393
hormones 191–197, 281–289
Hsp70 expression 379–384
hydraulic architecture 385–393, 510–521
Hypholoma fasciculare 153–161
Hypogastrura tullbergi 353–356

ideal free distribution 492–503
iguana 40–48
immunity levels 462–474
Impatiens glandulifera 530–539
incubation efficiency 305–312, 782–791
insects 313–325, 332–338, 484–491, 750–757
  ants 475–483, 773–781
  butterflies 484–491
  cockroaches 802–807
  leaf engineers 575–584
  spiders 734–741, 802–807
interaction effects 153–161, 640–648
irradiance 633–639, 682–689
isopods 596–607
isotope tracer 454–461

Juncus roemerianus 596–607
juvenile 73–78, 167–174, 462–474

kinematics 204–215
Koeleria cristata 758–765

Labridae 557–562
Lacerta oxycephala 842–850
Lacerta vivipara 135–140
Larus fuscus 445–453
Larus hyperboreus 617–622
Larus ridibundus 281–289
larval growth 640–648
latitudinal 167–174
layered substrate 664–670
leaf 403–412, 633–639, 682–689
  CO_2 exchange 49–57, 808–816
  miners 575–584

© 2002 British Ecological Society,

leaf area index 49–57, 727–733
Lepidoptera 484–491
lichens 808–816
life history 135–140, 750–757
life span 313–325, 379–384
life tables 313–325, 835–841
light 258–267, 633–639, 826–834
  availability 433–444
  limitation 232–240, 326–331
Lindera umbellata 826–834
litter 585–595, 596–607, 727–733
lizards 135–140, 817–825, 842–850
local adaptation 128–134
locomotion 40–48, 817–825
Lolium perenne 198–203, 394–402
Loranthaceae 128–134, 657–663
lowland mires 67–72
Lycosa tarantula 734–741

Macaranga triloba 475–483
macroecology 419–432
Madagascar 714–726
Magnolia salicifolia 826–834
Malacostraca 357–366
mammals 58–66, 394–402, 504–509, 773–781
mark–recapture 462–474
mate-locating 484–491
maternal effects 273–280, 281–289
maternal investment 135–140, 742–749
maximal acceleration capacity 484–491
mean residence time, nitrogen 826–834
medflies 313–325
Mediterranean 27–39, 773–781
mesic trees 113–121, 510–521
metabolic cold adaptation 332–338
metabolic cost 241–251
metabolic rate 175–190, 332–338, 671–681
metabolism 40–48, 73–78, 633–639
metamorphosis 640–648
Microtus ochrogaster 58–66
midrib damage 575–584
minirhizotron 198–203
mistletoes 128–134, 657–663
mite 353–356
models 27–39, 106–112, 268–272, 413–418
  leaf area index 727–733
  multiple regression 714–726
Molinia caerulea 198–203
montane 510–521
Monte Carlo simulation 504–509
moulting 357–366
mutualism 252–257, 475–483
mycelial cords 153–161
mycorrhiza 346–352, 742–749, 758–765
Myotis evotis 18–26
Mytilus edulis 339–345

Nardus stricta 198–203, 664–670
Nauphoeta cinerea 802–807
Nerodia sipedon 766–772
net assimilation rate 682–689
niches 258–267, 690–695, 714–726
nitrate 258–267
nitrogen 4–17, 403–12, 433–44, 727–33
  limitation 563–74
  spatial variation 454–461
  transport 191–197
nitrogen-use efficiency 67–72, 826–834
Niveoscincus metallicus 817–825
non-indigenous species 530–539
nonlinear 413–418
Nothofagus solandri 657–663
nutrient 223–231, 475–483, 705–713
  heterogenity 664–670
  limitation 122–127, 326–331
  uptake 339–345, 454–461, 742–749
nutritional stress limitation 623–632

Onychiurus spp. 353–356
organochlorines 617–622
Oryctolagus cuniculus 394–402
Oryza sativa 727–733
oscillatory stroking 106–112
outcrossing rate 79–88
oxygen provision 241–251
oxygen consumption, postexercise 40–48
ozone 4–17, 792–801

palm leaves 808–816
Pancratium sickenbergeri 99–105
Pararge aegeria 484–491
parasite 128–134, 657–663, 750–757
parental care 175–90, 241–51, 671–81
  nest attendance 782–791
Parus spp. 851–857
path analysis 682–689, 742–749
penguins 141–148, 175–190, 671–681
Peraxilla tetrapetala 657–663
perennials 79–88, 198–203
performance measurements 273–280
Periplaneta spp. 802–807
pH, *Carex* spp. and mire 67–72
Phallus impudicus 153–161
Phanerochaete velutina 153–161
phenolics 122–127, 216–222, 575–584, 596–607, 649–656
phenotype plasticity 510–521, 640–648, 835–841
Phleum pratense 454–461
phloem 575–584
phosphorus 346–352, 403–412, 758–765
photo-acclimation 808–816
photosynthesis 232–240, 727–733, 808–816
physical dormancy 252–257
physiological aggressiveness 167–174
physiological integration 191–197
pigment 445–453
Pinus 385–393
Pinus palustris 113–121, 596–607
Pinus ponderosa 510–521
pioneer species 258–267
Piper spp. 258–267
Pithecoctenium crucigerum 433–444
plant attributes 296–304
plant functional types 4–17
plant metabolism 633–639
plant strategies 326–331, 563–574
plant traits 585–595
plant water relationships 367–378
plant–plant communication 128–134
plant–pollinator interaction 492–503
plant–soil interactions 367–378
plasticity 67–72
Plicosepalus acaciae 128–134
Podarcis melisellensis 842–850
pollination 413–418
Populus tremuloides 792–801
prey 204–215, 766–772, 802–807, 842–850
principal components analysis 403–412
production 563–574
productivity, *Carex* spp. 67–72
protein 313–325
Pseudoroegneria spicata 758–765
Pygoscelis spp. 141–8, 175–90

quasi-steady model 106–112
Quercus spp. 113–21
Quercus virginiana 596–607

© 2002 British Ecological Society,

rainfall 403–412
ramets 191–197
ranging behaviour 504–509
red:far red ratio 258–267
reindeer herbivory 649–656
relative growth rate 223–231, 682–689
  Asteraceae 122–127
remote sensing 851–857
repeatability 273–280
reproduction 18–26, 289–295, 413–418
  costs 817–825
  insects 313–325, 734–741
reproductive output 623–632, 835–841
reproductive trade-off 705–713
reptiles 40–48, 135–140, 817–825
  snakes 204–215, 766–772
resource 191–197, 545–556
  allocation 73–78, 484–491, 705–713, 802–807
  availability 475–483, 714–726
respiration 27–39, 162–166, 596–607
response traits 545–556, 585–595
retrospective growth analysis 649–656
roads 504–509
rodent seed predation 773–781
roosts, natural 18–26
root 198–203, 223–231, 454–461, 664–670
  fine 113–121, 696–703
  life span 113–121, 198–203, 326–331
root parasite 128–134

Salix polaris 649–656
Salmo salar 73–78
Salmo trutta 167–174
salt marsh 596–607
sandhill 113–121
sap flow 696–703
saprotrophic basidiomycetes 153–161
Sarcobatus vermiculatus 367–378
scale 346–352, 545–556
sclerophylls 403–412, 633–639
scrubland 773–781
secondary compounds 122–127
sedges 4–17, 67–72
seed 296–304, 608–616, 742–749
  predation 252–257, 773–781
seedling 216–222, 223–231, 530–539
  establishment 433–444, 608–616
  relative growth rate 122–127
self-pollination 79–88
self-shading 232–240
senescence 727–733, 734–741
sex allocation 623–632
sex ratio 649–656
sexual dimorphism 766–772, 842–850
sexual selection 89–98
shade 222–231, 433–444, 808–816
shells 339–345, 714–726
shrub 4–17, 216–223, 232–240, 367–378
  evergreen 403–412, 664–670
Simpson's paradox 346–352
size 419–432, 492–503, 640–648, 835–841
  clutch 305–312, 782–791
  prey 766–772, 842–850
skewed distribution 419–432
skink 817–825
snail 623–632
snakes 204–215, 766–772
soil 162–166, 367–378, 433–444, 664–670
soil water respiration 27–39
Sorbus aucuparia 608–616
Sorghum bicolor 727–733
speciation 419–432
species-level response 346–352
specific leaf area 216–222, 682–689
specific root length 113–121
sperm competition 623–632
spider 734–741, 802–807
spinescence 216–222
standard metabolic rate 167–174
Storeria dekayi 766–772
structure, tree 385–393
survival 357–366, 462–470, 530–539
Svalbard 353–356
swimming costs 106–112, 141–148, 175–190
swimming performance 290–295, 557–562
Synchaeta pectinata 835–841
synchrony 413–418

Tabebuia rosea 433–444
tail loss 817–825
tannins 649–656
temperature 27–39, 175–190, 258–267, 290–295
  and eggs 305–312, 782–791, 835–841
  and growth 503–539, 640–648
territorial predator 734–741
territory quality 851–857
testosterone 281–289
Tetracanthella arctica 353–356
Tetradymia glabrata 367–378
Thamnophis sirtalis 766–772
Thamnophis spp. 204–215
thermoregulation 18–26
tolerance 99–105
torpor 18–26
trade-offs 99–105, 545–556, 682–689, 705–713, 782–791
transponders 141–148
tree 385–393, 475–483, 575–584, 608–616
  and competition 792–801
  roots 113–21, 223–31
Trifolium pratense 454–461
tropical 403–412, 475–483
  seedlings 223–231, 258–267, 433–444
Turdus spp. 608–616

Uca pugilator 89–98
ultradian activity rhythm 58–66
understorey species 585–595, 826–834
unpredictability 690–695
UV-B radiation 4–17

variable environment 313–25
variation, spatial 773–781
vegetation response 585–95
velocity assay 268–272
venom optimization theory 802–807
visual fields 522–529
vole 58–66

water absorption 696–703
water availability 79–88
water potential 258–267, 385–393
weaning mass 462–474
wind 826–834
wing propulsion 106–112
wood decay 153–161
woodland 153–161, 851–857
woody species 223–231

xeric trees 113–121, 510–521
xylem cavitation 367–378

yolk 135–140, 281–289, 445–453

zooplankton 835–841

© 2002 British Ecological Society,